


Saline-Filled Testicular Prosthesis



The cosmetic benefits provided by a testicular prosthesis have been reported by numerous investigators to result in psychological benefits and high levels of patient satisfaction.¹





Saline-Filled Testicular Prosthesis

Suture tab enables physicians to secure the device in a set position, if desired, eliminating unwanted implant movement within the scrotum.

Self-sealing injection port allows for filling the device with sterile saline solution.



Ordering Information

Coloplast Interventional Urology Surgical Support 800-258-3476

Product	Size	Α	В	Order Number
Torosa	Extra Small	2.2 cm	3.0 cm	450-1323
	Small	2.5 cm	3.5 cm	450-1325
	Medium	2.7 cm	4.0 cm	450-1327
	Large	2.9 cm	4.5 cm	450-1329

including educational videos, filling instructions, coding guide and additional ordering information, visit **ColoplastMD.com** or scan the QR Code below.

For additional Torosa resources



Multiple sizes are available to meet the needs of adult, adolescent and pediatric patients.

References

- 1. Clinical data on file at Coloplast Corp. and within the device labeling.
- 2. Turek, Paul J. et al. Safety and Effectiveness of a New Saline Filled Testicular Prosthesis. J Urol. 2004; 172:1427 143.
- 3. Hayon S, Michael J, Coward R M. The Modern Testicular Prosthesis: Patient Selection and Counseling, Surgical Technique, and Outcomes. Asian J Androl 2020; 22:64-9.

TOROSA® SALINE-FILLED TESTICULAR PROSTHESIS BRIEF STATEMENT

Indications: The Coloplast TOROSA Saline-Filled Testicular Prosthesis is intended for use when cosmetic testicular replacement is indicated i.e., in the case of agenesis or following the surgical removal of a testicle. Contraindications: The implantation of testicular prostheses is contraindicated in the presence of infection or untreated neoplasm.

Warnings: This device contains solid silicone elastomer. The risks and benefits of implanting this device in patients with lupus (e.g., SLE or DLE), scleroderma (e.g., progressive systemic sclerosis), myasthenia gravis, or documented sensitivity to silicone should be carefully considered. The issue of the possible relationship between silicone and various diseases has been and continues to be the subject of scientific and medical debate. Sepsis or hemorrhage may result from the placement of any foreign object in the body.

Excessive fibrous capsular formation or contracture may occur around any implant placed in contact with soft tissues.

Precautions: Each prosthesis should be checked for patency prior to surgery and continuously monitored throughout the surgical procedure to ensure that the structural integrity of the implant is not compromised in any way.

The action of drugs (such as antimicrobials, chemotherapy agents or steroids) in contact with the prosthesis has not been tested by the manufacturer, and their use cannot be recommended. Each physician who chooses to use drugs in combination with this prosthesis must assure compatibility of the drug with silicone elastomer.

A thorough preoperative consultation should include a discussion between the patient and physician of all available treatment options and their risks and benefits.

 $\label{potential} \textbf{Potential Complications:} \ \ \textbf{Potential complications include pain and discomfort.}$

The information provided is not comprehensive with regard to product risks. For a comprehensive listing of indications, contraindications, warnings, precautions, and adverse events refer to this product's Instructions for Use. Alternatively, you may contact a Coloplast Representative at 1-800-258-3476 and/or visit the company website at www.coloplast.us.

Caution: Federal law (USA) restricts this device to sale by or on the order of a physician.

